









Math 246 Review 2:

1. Draw a bar diagram to show how to solve each of the following word problems, and write the associated equations:

- Addition and subtraction bar diagrams need labels
- Multiplicative comparison bar diagrams need labels
- Other multiplication and division bar diagrams do not need labels
- Each diagram needs an addition or multiplication equation (which may be an unknown part equation)
- Some bar diagrams should also have a subtraction or division equation.

<p>a. Sandra had 4 erasers left after she gave 7 erasers to her friends. How many erasers did she start with?</p>  <p>$7+4=?$ gave has left optional: $?-7=4$</p>	<p>b. Marie has 8 blue balloons and 5 red balloons. How many more red balloons than blue balloons does she have?</p>  <p>$8-5=?$ $5+?=8$</p>
<p>c. Kyle has 9 transformers. He has 3 more than his sister. How many does his sister have?</p>  <p>$9-3=?$ $3+?=9$</p>	<p>d. Amanda has 6 origami cranes. How many more does she need to make to have 10 origami cranes?</p>  <p>$10-6=?$ has needs to make $6+?=10$</p>
<p>e. Paul had 14 cookies. He gave some to his brother, and now he has 8. How many cookies did he give to his brother?</p>  <p>$14-8=?$ $8+?=14$</p>	<p>f. A tootsie roll costs 4¢. Ross has 24¢. How many tootsie rolls can he buy?</p>  <p>$24 \div 4 = ?$ $4 \times ? = 24$</p>
<p>g. A toy train can go 20 feet in 5 seconds. How many feet can it go in one second?</p>  <p>$20 \div 5 = ?$ $5 \times ? = 20$</p>	<p>h. A Jar of jam has 8 ounces of jam in it. How many ounces of jam are in 5 jars?</p>  <p>$8 \times 5 = ?$</p>

g. A toy train can go 20 feet in 5 seconds. How many feet can it go in one second?

$20 \div 5 = ?$
 $5 \times ? = 20$

h. A Jar of jam has 8 ounces of jam in it. How many ounces of jam are in 5 jars?

$8 \times 5 = ?$

i. John has 4 pencils. Nathan has 5 times as many pencils as John. How many pencils does Nathan have?

$4 \times 5 = ?$

j. Kyle has 24 crayons. He has 3 times as many crayons as Clara. How many crayons does Clara have?

$24 \div 3 = ?$
 $3 \times ? = 24$



2. Show **two ways** of doing each calculation that are **different from the standard algorithm**. Know how to show at least one strategy for each on an open number line.

a. $36 + 29$

c. $92 - 38$

There are many correct solutions. For example:


a. open number line solutions:



a. other solutions:

$30 + 20 = 50$ $6 + 9 = 15$ $50 + 15 = 65$	$36 + 29 = 35 + 30 = 65$
--	--------------------------

c. open number line solution:



c. other solutions

$90 - 30 = 60$ $60 - 8 = 52$ $52 + 2 = 54$	$90 - 30 = 60$ $2 - 8 = -6$ $60 - 6 = 54$	$92 - 38 = 94 - 40 = 54$
--	---	--------------------------

3. Explain (using appropriate base 10 language) the following two steps in the standard subtraction algorithm:

$$\begin{array}{r}
 628 \\
 - 293 \\
 \hline
 5
 \end{array}
 \Rightarrow
 \begin{array}{r}
 512 \\
 \cancel{6} \cancel{2} 8 \\
 - 293 \\
 \hline
 5
 \end{array}
 \Rightarrow
 \begin{array}{r}
 512 \\
 \cancel{6} \cancel{2} 8 \\
 - 293 \\
 \hline
 35
 \end{array}$$

In the first step, I **trade a hundred for 10 tens**. Write down that there are now 5 **hundreds** (cross out 6), and there are now **12 tens** (instead of 2).

In the second step, I take away 9 **tens** from 12 **tens**, which leaves 3 **tens**. Write 3 in the tens place of the answer.

4. Show how to solve each of these using the appropriate expanded algorithm:

- a. $478 + 394$ b. $723 - 186$ c. 246×87

$$\begin{array}{r}
 478 \\
 + 394 \\
 \hline
 12 \\
 160 \\
 1200 \\
 \hline
 1372 = 537
 \end{array}
 \quad
 \begin{array}{r}
 600 \\
 \cancel{7} \cancel{0} \cancel{0} \\
 - 100 \\
 \hline
 500
 \end{array}
 +
 \begin{array}{r}
 110 \\
 \cancel{1} \cancel{1} \cancel{0} \\
 - 80 \\
 \hline
 30
 \end{array}
 +
 \begin{array}{r}
 13 \\
 \cancel{1} \cancel{3} \\
 - 6 \\
 \hline
 7
 \end{array}$$

$$\begin{array}{r}
 246 \\
 \times 87 \\
 \hline
 142 \\
 1280 \\
 1400 \\
 480 \\
 13200 \\
 16000 \\
 \hline
 21322
 \end{array}$$

$$\begin{array}{r}
 673R5 \\
 12 \overline{)8081} \\
 \underline{4800} \quad 400 \\
 3281 \\
 \underline{2400} \quad 200 \\
 881 \\
 \underline{480} \quad 40 \\
 401 \\
 \underline{360} \quad 30 \\
 41 \\
 \underline{36} \quad 3 \\
 5 \quad 673
 \end{array}$$

5. Show how to solve the following problem using scaffolding division in a way that uses easier multiplication facts than the most efficient solution:




$8081 \div 12$



6. a. Explain how knowing the commutative law of multiplication helps children learn the multiplication facts

It means that if a student knows one multiplication fact already (6x5), then they don't have to learn the turn-around fact (5x6) if they know the commutative law. It also lets them choose between two strategies/interpretations (5 groups of 6 or 6 groups of 5) to choose the one that they know best how to find the product.

b. Draw a diagram and write an (in words) explanation that shows why the commutative law of multiplication makes sense (you may show it for a specific example)




<p>version 1: Commutative law: $axb = bxa$ $6 \times 5 = 5 \times 6$</p>  <p style="text-align: center;">$6 \times 5 = 6$ groups of 5</p> <p style="text-align: center;">$5 \times 6 =$ 5 groups of 6</p> <p>6 groups of 5 and 5 groups of 6 are two ways of counting the same squares, so they are equal.</p>	<p>version 2: Commutative law: $axb = bxa$ $6 \times 5 = 5 \times 6$</p>  <p style="text-align: center;">$6 \times 5 =$ 6 groups of 5</p>  <p style="text-align: center;">$5 \times 6 =$ 5 groups of 6</p> <p>6×5 and 5×6 are equal because the arrays have the same length and width so they are the same size and shape</p>
---	--

7. a. Explain how knowing the distributive law of multiplication over addition helps children learn the multiplication facts

Children can break an unknown multiplication fact down into two known multiplication facts and add them together to find the total product

b. Draw a diagram and write an (in words) explanation that shows why the commutative law of multiplication makes sense (you may show it for a specific example).

There are 3 correct explanations that I can think of:

<p>version 1:</p> <p>Distributive Law $a(b+c)=ab+ac$ $3x(5+2)=3x5+3x2$</p>  <p>The array for $3x(5+2)$ can be split into two smaller arrays: $3x5$ and $3x2$, so the number of squares in $3x(5+2)$ is the same as the squares in $3x5$+ the squares in $3x2$.</p>	<p>version 2:</p> <p>Distributive Law $a(b+c)=ab+ac$ $3x(5+2)=3x5+3x2$</p>  <p>3 groups of 7 is the same amount as 3 groups of 5 and 3 groups of 2 because each 7 can be split into 5 and 2</p>	<p>version 3:</p> <p>Distributive Law $a(b+c)=ab+ac$ $3x(5+2)=3x5+3x2$</p>  <p>7 groups of 3 is the same as 5 groups of 3 and 2 more groups of 3</p>
---	---	---

8. Write **a.** a partition and **b.** a measurement division word problem for $36 \div 4$.


a. A partition division problem tells you the number of sets. For example “I have 36 cookies that I want to put equally into 4 bags. How many cookies should go in each bag?”

b. a measurement division problem tells you how many is in each set. For example: “I have 36 cookies. I want to put 4 cookies into each bag. How many bags can I fill?”

9. Write a word problem for 32×14

Example: A shirt costs \$32. How much do 14 shirts cost?


10. Show how to compute 438×49 using the lattice algorithm



11. a. Show how to compute $\begin{array}{r} 548 \\ \times 37 \\ \hline \end{array}$ using the standard algorithm.

<p>a. Compute with the standard algorithm</p> $\begin{array}{r} 1 \quad 2 \\ 3 \quad 5 \\ 548 \\ \times 37 \\ \hline 3836 \\ 16440 \\ \hline 20276 \end{array}$	<p>b. Before computing 3×8 in the standard algorithm we write a 0 in the partial product. Explain why we write a 0 there. <i>3 is 3 tens, so the product should go in the tens place (24 tens), so we put a 0 in the ones place.</i></p>	<p>c. When we compute $3 \times 8 = 24$ on the standard algorithm, we write 4 in the tens place, and we write 2 above the tens place. Why does 4 go in the tens place, and why does 2 go above the tens place? <i>The 3 is 3 tens, so 24 is 24 tens, so 4 should go in the tens place (because it's 4 tens). The 2 is 20 tens = 200, so it needs to get added in the next place value. I'll get the next place value (hundreds) when I multiply 30×40, so I should put 2 above the 4 to make it easy to add on.</i></p>
---	--	--

12. a. Sketch an array diagram for: 57×82

	<p>Notes about the array: One side must have length $80+2$ The other side must have length $50+7$ The array must be subdivided into 4 parts where the numbers split between tens and ones.</p> <p>The lengths do not have to be to scale or even approximately to scale unless I provide a grid.</p>
<p>expanded:</p> $\begin{array}{r} 82 \\ \times 57 \\ \hline 14 \\ 560 \\ 100 \\ 4000 \\ \hline 4674 \end{array}$	<p>standard:</p> $\begin{array}{r} 1 \\ 82 \\ \times 57 \\ \hline 574 \\ 4100 \\ \hline 4674 \end{array}$ <p>note that you could optionally choose to double color the 7 in 574 as red and purple, and the 1 in 4100 as blue and green.</p>

13. a. Write a division problem with a remainder where the answer that makes sense is the quotient
This needs to be a problem where it would make sense to discard the remainder: "I have 25 cookies. 4 cookies can fit in a bag. **How many bags can I fill?**"

b. Write a division problem with a remainder where the answer that makes sense is the quotient+1
This needs to be a problem where you have to keep and make a group for the remainder: "There are 25 children. 4 children can ride in each car. How many cars do we need **to take all of the children** to the zoo?"

(14. Analyze and explain an error pattern or an alternate algorithm for addition, subtraction, multiplication or division.—see homework for examples)

15. Show two ways of figuring out 4×9 using efficient strategies.

An efficient strategy is faster than skip counting ($4+4+\dots+4$), but is not just a memorized answer.


<p>Focusing on the 4, and thinking of the problem as 4 groups of 9:</p> <p>4×9</p> <p>Double twice:</p> <p>$9 \times 2 = 18$</p> <p>$18 \times 2 = 36$</p>	<p>Focusing on the 9, and thinking of the problem as 9 groups of 4:</p> <p>$4 \times 10 = 40$</p> <p>(9 groups of 4)</p> <p>$4 \times 10 = 40 - 4$</p> <p>$= 36$</p> <p>(10 groups of 4 - 1 group of 4)</p> <p>or use the 9's pattern: 4×9 will have 3 as the tens digit (one less than 4). The tens and ones digits will add to 9: $3 + \underline{6} = 9$ so 36.</p>
--	--

16. Show two ways of figuring out 6×7 using efficient strategies.

<p>Thinking 6 groups of 7, and breaking 6 into 5 and 1:</p> <p>$6 \times 7 =$</p> <p>Six sevens</p> <p>$5 \times 7 = 35$</p> <p>five sevens</p> <p>$35 + 7 = 42$</p> <p>5 + 1 sevens</p>	<p>6 groups of 7 and $6 = 3 + 3$</p> <p>$3 \times 7 + 3 \times 7$</p> <p>$21 + 21$</p> <p>42</p>	<p>7 groups of 6 and $7 = 5 + 2$</p> <p>6×7</p> <p>Seven sixes</p> <p>five sixes = $5 \times 6 = 30$</p> <p>+ 2 sixes = $\frac{12}{42}$</p>
---	--	--


17. a. Show what a direct modeling type picture (so you could count each object to find the answer) of a partitive division solution for $19 \div 5$ would look like.

partitive means 5 is the number of sets



b. Show what a direct modeling type picture (so you could count each object to find the answer) of a **partitive** measurement division solution for $19 \div 5$ would look like. (one of these should have been measurement).

measurement means the 5 is the size of each set



18. Solve by long division $4793 \div 13$

$$\begin{array}{r}
 368 \text{ R}9 \\
 13 \overline{)4793} \\
 \underline{39} \\
 89 \\
 \underline{78} \\
 113 \\
 \underline{104} \\
 9
 \end{array}$$